



Review Article

The Short- and Long-Term Clinical, Radiological and Functional Consequences of COVID-19



Yang Gao^{a,1}, Wei-quan Liang^{b,1}, Yi-ran Li^{b,1}, Jian-xing He^{c,d,*}, Wei-jie Guan^{b,c,d,**}

^a Department of Pulmonary and Critical Care Medicine, Beijing Anzhen Hospital, Capital Medical University, Beijing, China

^b Department of Respiratory and Critical Care Medicine, Foshan Second People's Hospital, Affiliated Foshan Hospital of Southern Medical University, Foshan, Guangdong, China

^c State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, Guangzhou Institute of Respiratory Health, First Affiliated Hospital of Guangzhou Medical University, Guangzhou Medical University, Guangzhou, China

^d Department of Thoracic Surgery, Guangzhou Institute for Respiratory Health, The First Affiliated Hospital of Guangzhou Medical University, China

ARTICLE INFO

Article history:

Received 22 February 2022

Accepted 4 March 2022

Available online 13 April 2022

Keywords:

COVID-19

Long COVID

Recovery

Symptom

Intervention

Radiology

Psychology

ABSTRACT

As with the rapid increase of the number of patients who have recovered from COVID-19 globally, there needs to be a major shift of the focus from rapid pathogen detection, treatment and prevention to the promotion of better recovery. Notwithstanding the scarcity of our understandings, recent studies have unraveled a plethora of pulmonary and systemic consequences which require medical attention. These consequences remained as of the end of follow-up which ranged from 1 month to 1 year. Here, we review the consequences of COVID-19 in terms of the residual symptoms, radiological and functional manifestations, and identify the potential risk factors that contribute to a prolonged recovery. We also summarize the benefits of clinical interventions (particularly the pulmonary rehabilitation program), and address several undetermined concerns and key future research directions.

© 2022 Published by Elsevier España, S.L.U. on behalf of SEPAR.

Consecuencias clínicas, radiológicas y funcionales a corto y largo plazo de la COVID-19

RESUMEN

Como consecuencia del rápido aumento del número de pacientes que se han recuperado de la COVID-19 en todo el mundo, es necesario cambiar el enfoque de la detección rápida del patógeno, el tratamiento y la prevención para promover una mejor recuperación. A pesar de la escasez de nuestros conocimientos, estudios recientes han desvelado una pléthora de consecuencias pulmonares y sistémicas que requieren atención médica. Estas consecuencias se mantienen al final del seguimiento, que oscila entre 1 mes y 1 año. Aquí se hace una revisión de las consecuencias de la COVID-19 en términos de síntomas residuales y manifestaciones radiológicas y funcionales y se identifican los posibles factores de riesgo que contribuyen a una recuperación demorada. También se resumen los beneficios de las intervenciones clínicas (en particular el programa de rehabilitación pulmonar) y se abordan varias preocupaciones no resueltas y direcciones clave de investigación futura.

© 2022 Publicado por Elsevier España, S.L.U. en nombre de SEPAR.

Palabras clave:

COVID-19

COVID persistente

Recuperación

Síntomas

Intervención

Radiología

Psicología

* Corresponding author.

** Corresponding author.

E-mail addresses: hejx@vip.163.com (J.-x. He), battery203@163.com (W.-j. Guan).

¹ Yang Gao, Wei-quan Liang, and Yi-ran Li contributed equally to the work. Jian-xing He and Wei-jie Guan contributed equally to the work.

Introduction

Coronavirus disease of 2019 (COVID-19) has strained the medical infrastructure, personnel and resources worldwide. The burden continues to increase amid the emergence of new variants and the progressively waning immunity after global vaccine roll-out. According to the latest statistics, the total number of laboratory-confirmed cases of COVID-19 has exceeded 0.4 billion globally,¹ despite the dramatically improved rate of survival.

Although a certain number of patients ultimately developed severe or critical illness, symptoms gradually resolved among most patients at convalescence.^{2–4} The number of patients who recover from COVID-19 will continue to increase considerably in the next foreseeable future. This constitutes a pressing need for caring a massive population which has not been fully characterized previously. Not all patients achieved complete remission of symptoms, radiologic findings and pulmonary function despite a prolonged convalescence phase.^{2,3,5–11} The consequences contributed to the ongoing suffering from physiologic dysfunction and the impaired quality-of-life which markedly dampened the overall well-being. Given the overwhelming tasks of disease prevention and control, global research efforts have characterized the clinical presentations at disease onset or during hospital admission, to identify the prognostic factors and disease phenotypes, to design and validate rapid diagnostic tests, and to verify the effects of therapeutic interventions and vaccines. Recent studies have documented the manifestations of consequences which ranged from respiratory to systemic (including psychological) disorders,¹² with the follow-up duration ranging from 3 months to 1 year.^{2–5,10,11,13–17} In light of the variable study designs and research focus, there remains a considerable knowledge gap in the understanding of the magnitude and heterogeneity of the consequences of COVID-19, particularly in the era when the medical focus has been shifted from the diagnosis and treatment to the rehabilitation.

Exploring the spectrum of consequences among patients recovered from COVID-19 might help characterize the residual disease burden and identify the major predictors of adverse outcomes, which have important implications to the early diagnosis and accelerate the development of novel interventions. Here, we review the major manifestations of short- and long-term consequences, highlighting the clinical characteristics and predictors of having any consequences. We also summarize the latest clinical interventions and elaborate the important future research directions.

Clinical characteristics at acute phase

Airborne droplets or virus-laden materials were the dominant sources of infection.¹⁸ After acquisition of virus, common symptoms which consisted of fever, cough, fatigue, dyspnea, sore throat and headache typically occurred after the median incubation period of 4 days.¹⁹ However, >50% of patients remained afebrile on hospital admissions and developed fever thereafter. Gastrointestinal symptoms, such as loss of appetite, nausea and vomiting, were less common as compared with those in patients with avian influenza.²⁰ Neurological manifestations (e.g., myalgia, encephalopathy, dysgeusia and anosmia) were specific to aid in the diagnosis although appearing less common.²¹ Few patients developed autoimmune disorders, including haemolytic anemia,²² immune thrombocytopenia,²³ Guillain-Barre syndrome,²⁴ and Kawasaki disease-like syndrome in children.²⁵

On laboratory examination, the most common manifestations consisted of lymphopenia, elevated C-reactive protein and elevated liver enzymes.¹⁹ These manifestations have important clinical implications for prognosis. Lymphopenia (seen in ~85% of patients), has been associated with the risk of progression to critical illness and death.¹⁹ Several radiologic manifestations on chest X-ray or

computed tomography (CT), such as ground-glass opacity, pulmonary infiltration, and consolidation usually developed at later stages of progression.²⁶ The radiologic abnormality, albeit rapidly evolving during progression,²⁶ could be readily detected at early stages among most patients and continued to persist.¹⁹ Lung function findings regarding airflow limitation and lung volumes were usually unremarkable.²⁷ By contrast, diffusing capacity impairment and reduced exercise tolerance were more specific to reflect functional impairment of COVID-19.²⁸

The path of disease progression

Although approximately 85% of patients developed mild diseases,¹⁹ the risk of progression to critical illness (e.g., severe pneumonia, acute respiratory distress syndrome, sepsis, multisystem organ failure) or death among the remaining patients could not be underestimated. Older age, greater symptom burdens, severe hypoxemia,²⁹ and comorbidities (e.g., hypertension, diabetes, heart disease, chronic obstructive pulmonary disease) have been associated with a worse outcome.³⁰ Furthermore, certain chest radiologic³¹ and laboratory parameters (e.g., decreased CD3+, CD4+, and CD8+ T cell count, higher neutrophil-to-lymphocyte and neutrophil-to-CD8+ T cell ratio) might provide prognostic implications.³² Prevention of progression to critical illness or death has been the core mission of the clinical management, because the severity of illness correlated negatively with the likelihood of, and the time needed for, clinical recovery.³³

Short- and long-term consequences

Although the symptoms or signs of COVID-19 ultimately resolved among the majority of patients, some suffered from persistent or new-onset symptoms and conditions which have been described as “long COVID”.³⁴ The definition is continuously evolving, although a new consensus definition for “post COVID-19 condition” has recently been proposed.³⁵ A commonly accepted concept refers long COVID to the symptoms that continue or develop after acute COVID-19 infection which cannot be explained by an alternative diagnosis.³⁶ However, this could be seen in many COVID-19 survivors >6 months after recovery.³⁷ Thus, we summarize the short- (4 to 12 weeks after disease onset) and long-term (12 weeks to 1 year or longer) consequences of COVID-19 according to the ongoing symptomatic and post COVID-19 syndrome stages below, as proposed by the UK National Institute for Health and Care Excellence guideline.³⁶

Short-term consequences

Symptoms and signs: At 4–12 weeks after the initial symptom onset, a considerable proportion of patients reported residual symptoms such as cough (2%–40.3%), dyspnea (5.5%–91.5%), chest pain (0.2%–42%), sputum production (<10%), cognitive impairment (17%–28.3%), and fatigue (17%–84.8%), while fever might persist for a couple of weeks without secondary infection (Table 1).^{37–40} Over 60% of patients reported amelioration of olfactory and/or gustatory dysfunction within the first 2 months (with the percentage decreasing to 40% and 20%, respectively), followed by a slower rate of recovery thereafter.⁴¹ Psychological disorders (e.g., post-traumatic stress disorder [PTSD]) occurred in 24.5% of survivors at 1 month after discharge and in 22.4% of patients at 3 month follow-up.⁴² Insomnia, anxiety and depression could still be identified at 3 months.⁴²

Chest radiologic findings: Chest radiologic abnormalities could be identified in >75% of patients at initial presentation¹⁹ and could persist at convalescence. Ground-glass opacification (GGO) was

Table 1

Proportion of patients with persistent symptoms at the short and long terms.

Persistent symptoms	4-12 weeks	6 months	1 year or longer
<i>Patients with at least 1 symptom</i>	41.0-94.0%	37.0-81.0%	45.0-60.9%
<i>Common symptoms</i>			
Fatigue	17.0-84.8%	8.0-85.0%	10.0-60.9%
Dyspnea	5.5-91.5%	11.9-42%	2.7-48%
Chest pain	0.2-42%	0-21.0%	0-15.8%
Cough	2.0-40.3%	2.1-24.0%	3.0-29.0%
Hair loss	13.3-28.6%	2.5-26.3%	11.0-36.2%
Anxiety or depression	4.3-45.1%	17.39-26.7%	3.3-42.0%
Sleep disorder (e.g. insomnia)	3.6-53.6%	5.42-35%	10.7-43.3%
Impaired memory or poor concentration	8.1-41.0% (up to several months)		
<i>Less common symptoms</i>			
Joint pain	7.6-27.3%	9.0-33.0%	12.8-32.5%
Headache	1.6-61.0%	2.0-26.0%	2.3-17.6%
Myalgia	4.5-26.0%	2.0-38.0%	4.0-27.0%
Anosmia	1.7-10.0%	3.3-18.0%	1.3-26.3%
Ageusia	2.5-10.0%	3.0-13.0%	1.4-30.2%
Post-traumatic syndrome disorders	14.2-28.0% (up to several months)		
Sputum production, weight loss, sweating, skin rash, sore throat, edema of lower limbs, palpitation, diarrhea, constipation, nausea, decreased appetite, abdominal pain, dizziness, anorexia	<10% (up to several months)		
Fever	<5% (up to several weeks)		

Abnormalities reported within 4-12 weeks^{53,75,99-104}Abnormalities reported at 6 months^{2,3,63,70,74,83,100,105-107}Abnormalities reported at 12 months or longer^{2,46,61,62,64-67,72,108}

highly prevalent at discharge (99.5%) but was only detected among 37.3% of patients at 3 months. However, consolidation and crazy-paving pattern might have resolved at hospital discharge. At 1 month after discharge, GGO (42.4%, 37.5%) and reticular pattern (22.0%, 47.5%) were most common in patients with mild and severe COVID-19. These figures decreased to 29.7% and 28.3% for GGO and 13.0% and 36.7% for reticular patterns at 3 months.⁴³ By contrast, parenchymal bands, thickening of adjacent pleura and reticular lesions persisted in 34.4%, 13.9%, and 14.8% of patients at 3 months.⁴⁴

Lung function abnormality: Although occurring in a small proportion of patients during acute phase, pulmonary function remained abnormal in the early post-acute phase especially in patients with severe COVID-19.^{45,46} Reduced diffusing capacity, seen in 10-20% of patients at presentation, could substantially improve at 3 month in patients with severe COVID-19.⁴⁶⁻⁴⁸

Systemic complications: COVID-19 conferred notable adverse impacts on multiple organ systems, whose function was progressively normalized in the short term. Cardiovascular complications included myocarditis, myocardial infarction, heart failure, cardiogenic shock and arrhythmia.⁴⁹ COVID-19 appeared to be associated with persistent cardiac injury after recovery, particularly subclinical myocardial injury in the short term and diastolic dysfunction in the long term.⁵⁰ Despite the high incidence at acute phase,¹⁹ acute kidney injury could be generally resolved within 3 weeks without renal replacement therapy among patients with non-severe COVID-19;^{51,52} new-onset chronic kidney disease was unusual but could relapse after discharge.^{3,53} Acute liver injury or hepatitis occurred in few patients only and resolved during the acute phase among patients with non-critical illness, with minor impacts on long-term outcomes.^{30,54,55} Thyroid function might be impaired at acute phase in approximately 13.5% of patients and could be normalized within 2 months.⁵⁶ Hypercoagulability (e.g., venous and arterial thromboses) was common at acute phase, especially in patients with severe or critical illness,⁵⁷ but the incidence of venous thromboembolism fell to below 5% in the post-acute phase.³⁸

Short-term mortality: Although most patients with COVID-19 had asymptomatic or mild-to-moderate illnesses, findings from the initial outbreak indicated the fatality rate of approximately 1.4% in mainland China.¹⁹ Findings from Spain have also lent support to these earlier observations. In a multicenter Spanish

cohort of 12,126 patients, the mortality rate was 12.5% for non-hospitalized patients, 29.8% in hospitalized patients and 38.8% in patients admitted to the intensive care unit.⁵⁸ The key predictors of short-term mortality consisted of the age over 50, obesity, cardiac diseases, fever, dyspnea, lung infiltration, lymphopenia, D-dimer levels above 1000 ng/mL, and requirement of mechanical ventilation.⁵⁸ When taking into account the impact of the underlying diseases, in particular the chronic respiratory diseases, another multicenter Spanish cohort study of 5847 hospitalized patients with COVID-19 revealed a significantly higher 30-day mortality rate in patients with prior lung diseases than those without (29.5% vs. 17.9%).⁵⁹ The study also revealed a more impressive 30-day mortality rate (approximately 40%) in patients with pre-existing COPD.⁵⁹ Both studies, however, concurred that a greater age and higher D-dimer levels were associated with a significantly elevated risk of short-term death.^{58,59} Patients aged greater than 60 years were at a particularly high risk of death at the short term, with the overall in-hospital mortality rate of 23.5% in a retrospective single-center study in Spain.⁶⁰ Among these elderly patients, those with malignancy, chronic liver diseases, obesity and diabetes had an increased mortality risk despite the adjustment with the age.⁶² These studies have provided the scientific rationale for rapid identification of patients at risk of death and the implementation of more intensive therapeutic management.

Medium- to long-term consequences

Symptoms and signs: At week 12 and thereafter, symptoms or signs progressively waned but could still fluctuate over time. Some patients reported residual symptoms for up to 6 months or longer, especially those with severe COVID-19.^{3,61} Cough lasted for 6 months or longer in 20% of patients (Table 1),⁶² which might be associated with pulmonary fibrotic changes.⁶³ Breathlessness could persist for up to 1 year.² Chest pain or tightness was reported in approximately 15.8% of patients over 1 year.⁶⁴ Systemic symptoms such as fatigue persisted for 12 months among 10-60.8% of patients,^{2,58,64-68} which might be associated with the accentuated systemic inflammatory response and psychological disorders.⁶⁸ Persistent taste and smell disorders have been reported at 6 months (7% and 11%, respectively),³ although recovered progressively thereafter.² Albeit less commonly reported, arthralgia

and myalgia could persist for over 6 months.^{2,69–71} Hair loss was reported in >20% of patients at 6 months while decreased to 11% at 1 year.² Neurological disorders (e.g. sleep disorder, headache, dizziness) persisted with minor improvement for 1 year.^{2,62,71} Noticeably, 27% of patients suffered from sleep disorder at 6 months and 17% still had insomnia at 1 year.² Cognitive impairment was reported to be persistent, the magnitude of which fluctuated from 4 months to 1 year, and was more prevalent in patients with severe COVID-19.^{62,72–75} COVID-19 imposed a considerable burden on psychology. More than 20% of patients suffered from PTSD at 1 year or longer.⁷⁶ Anxiety and/or depression was reported in approximately 23% of patients at 6 months and 26% at 12 months.^{2,62,76}

Chest radiologic findings: During recovery, GGO and interstitial fibrotic changes remained common.^{2,44,53} More than 60% of patients with mild or moderate COVID-19 ultimately had resolved radiologic abnormalities. However, GGO and fibrotic changes could persist while consolidation was rare on chest CT at 1 year follow-up.^{2,77} Furthermore, the disease severity was associated with the magnitude of chest CT manifestations. At 3 months, pulmonary structural abnormalities were highly prevalent (up to 80.7%) in patients hospitalized in the intensive care unit due to ARDS.⁷⁸ Radiologic abnormalities persisted in 20% of patients with severe COVID-19 who did not require mechanical ventilation at 12 months.⁷⁹ Compared with patients who had completely resolved, a significantly higher proportion of patients diagnosed as having severe pneumonia and acute respiratory distress syndrome at the initial presentation had residual radiologic abnormality at 1 year.⁴⁴

Lung function abnormality: Restrictive and obstructive patterns and impaired diffusing capacity were common in COVID-19 survivors,⁸⁰ although improved considerably in non-critical survivors within 3 months.⁴⁷ The improvement of diffusing capacity, however, seemed to slow down at 6 months. For up to 12 months, impaired diffusing capacity was identified in 20-30% of patients who had moderate COVID-19, whose prevalence remained constant since month 6. By contrast, impaired diffusing capacity could be found in approximately 54% of patients with critical illness at 1 year.² In patients who had initially suffered from severe COVID-19 but did not require mechanical ventilation, impaired pulmonary function improved gradually over 1 year, with females having a higher risk of the ongoing impairment as compared with males.⁷⁹ Impaired diffusing capacity could persist for a longer period, for instance, up to 18 months after discharge, among 57.92% of survivors with more severe disease.^{46,81} Reduced exercise capacity was another important characteristic lung function change among survivors,³⁷ which improved gradually since month 3 in patients with severe COVID-19.⁷⁹

Systemic complications: As mentioned above, most systemic complications appeared to be transient. However, the estimated glomerular filtration rate decreased in 35% of patients after 6 months, 13% of which was categorized as the new-onset reduction of glomerular filtration rate despite the normal renal function at COVID-19 onset.³ At 6 months, hyperglycemia persisted in >30% of patients who experienced new-onset hyperglycemia during the hospitalization.⁸² A cohort study reported new-onset diabetes in 3.3% out of 1733 patients at 6 months.³




Figure 1. Acute, short- and long-term effects of COVID-19 on multiple organ systems. Clinical manifestations a patient might experience at a specific stage of COVID-19. eGFR: estimated glomerular filtration rate; PTSD: post-traumatic stress disorder.

Functional abnormality: The magnitude of improvement depended considerably on the initial disease severity. Most patients with severe illness reported a failure of recovery in physical activity at 6 months.⁸³ 45% of hospitalized survivors exhibited limitations on daily living activities at 7 months (Table 1).⁸⁴ The health care workers with severe COVID-19 showed improvement in functional fitness within 1 year, and rigorous intervention might be considered for accelerating the pulmonary rehabilitation (Fig. 1).⁸⁵

Medium- and long-term mortality: The adverse impact of COVID-19 on the mortality risk extended to the long terms. In a cross-sectional study conducted in France, the mortality rate was 62.5% during the intensive care unit stay and increased to 72.1% to month 6 among patients aged greater than 80 years who were receiving mechanical ventilation.⁸⁶ At 1 year, patients treated with corticosteroids had a markedly higher mortality rate as compared with those not (50.0% vs. 21.0%), although the data interpretation was limited by the small sample sizes.⁸⁷ A study conducted in Italy has also demonstrated a markedly higher mortality rate among hospitalized patients with COVID-19 than those without (48.4% vs. 33.9%) at 1 year.⁸⁸ The 1-year mortality rate was also higher in non-hospitalized emergency room visitors with COVID-19 than those without (18.0% vs. 8.7%).⁸⁸

Future research perspective

The continuously evolving global pandemic has elicited an unprecedented challenge for healthcare workers, policy makers and the public. Current research has outlined the clinical spectrum of the consequences, but this would surely benefit from the in-depth mechanistic insights which would unveil the biologic pathways that contribute to the ongoing disorders. Continuous profiling of the cytokine and chemokine expression levels within the lower airways (e.g., induced sputum) or peripheral blood will help understand the dynamics of residual inflammatory responses. The combination of chest imaging with high-resolution CT or functional magnetic resonance imaging with isotopes and lung function assessment (e.g., lung volume, diffusing capacity, exercise capacity) will provide clinicians an indispensable avenue for tracking the trend of changes in pulmonary functional abnormality.^{7–9,17,89} Further extension of the clinical observation is needed to reveal the long-term impact (e.g., 3–5 years) on symptom perception, lung function (particularly the exercise capacity), radiological characteristics, quality-of-life and psychology.^{10,11,16,90} This will help reveal the duration of recovery of individual symptoms which would normally take, and determine the optimal timing for evaluating the magnitude of recovery and defining complete or partial recovery.

Few intervention strategies have been proposed to accelerate recovery, with pulmonary rehabilitation demonstrating some promising findings.^{91,92} Anti-inflammatory medications or biologics might have a role in antagonizing the residual inflammation, especially when taking into account the pulmonary (e.g., cough, sputum production) and systemic manifestations (e.g., fatigue). There also lacks a crystal-clear indication for the population which would benefit most from therapeutic interventions after discharge. Whether non-pharmaceutical interventions are superior to drug therapy for improving the functional well-being (e.g., exercise capacity) merits further investigations from the personalized interventions. It would be important to address the optimal duration of such interventions should they be proven effective.

Previous studies have indicated a strong immune protective effect against acquiring re-infections among the COVID-19 rehabilitees.⁹³ The natural immunity has constituted the front line which prevents from being susceptible. However, approximately 4% of patients were susceptible to re-infections with the precise mechanisms not fully explored.^{93–95} Despite the formation of

pan-sarbecovirus antibodies,⁹⁶ it remains unclear whether COVID-19 survivors would be immune to other coronavirus. Continuous follow-up might help delineate whether re-infections were caused by the waning immunity alone, the underlying immunodeficiency, or emerging variants evading the established immunity. A minority of patients were tested continuously RNA-positive or re-positive after discharge which might not necessarily reflect prolonged viral shedding.^{97,98} A work-up with the government, epidemiologists and clinicians will help ascertain the true infectivity of COVID-19 survivors by gathering the information of the dynamic nucleic acid assays monitoring, enforcing symptom monitoring of patients and close contacts, and integrating multiple laboratory tests (e.g., viral culture, high-throughput sequencing). This will be important for redefining the community-based prevention measures, since no existing study has disclosed the infectivity of COVID-19 survivors who were discharged after having at least two consecutive RNA-negative laboratory test findings.

Conclusions

The world has come across a tough period when COVID-19 resulted in an overwhelming burden. Thanks to the efforts of the medical and scientific communities, the rapid deployment of effective vaccines and accelerated development of repurposed and novel medications, most patients have either partially or completely recovered from COVID-19. Although many questions remain unanswered, we keep optimistic that the ongoing efforts will better characterize the trajectory of clinical recovery and unveil new interventions which would promote recovery. The goal of enabling a better recovery would ultimately be realized for most COVID-19 rehabilitees.

Support statement

This work was supported by Guangzhou Institute for Respiratory Health Open Project (funded by China Evergrande Group) - Project No. 2020GIRHHMS09 and 2020GIRHHMS19 (Prof. Guan) and the Open Project of the State Key Laboratory of Respiratory Disease (SKLRD-OP-201909) (Dr. Gao).

Author contributions

Yang Gao, Wei-quan Liang, Yi-ran Li, Jian-xing He, and Wei-jie Guan drafted the manuscript; Wei-jie Guan, and Jian-xing He critically revised the manuscript. All authors have approved the final submission.

Conflicts of interest

The authors declared no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Acknowledgment

We thank Prof. Jiang Xie (Department of Pulmonary and Critical Care Medicine, Beijing Anzhen Hospital, Capital Medical University) for his helpful comments on the writing of the review.

References

- WHO Coronavirus (COVID-19) Dashboard. 2021. Accessed Feb 21st, 2022.
- Huang L, Yao Q, Gu X, Wang Q, Ren L, Wang Y, et al. 1-year outcomes in hospital survivors with COVID-19: a longitudinal cohort study. Lancet. 2021;398:747–58.
- Huang C, Huang L, Wang Y, Li X, Ren L, Gu X, et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. Lancet. 2021;397:220–32.

- Zhang J, Xu J, Zhou S, Wang C, Wang X, Zhang W, et al. The characteristics of 527 discharged COVID-19 patients undergoing long-term follow-up in China. *Int J Infect Dis.* 2021;104:685–92.
- Martinez-Garcia MA, Aksamit TR, Aliberti S. Bronchiectasis as a long-term consequence of SARS-CoV-19 pneumonia: future studies are needed. *Arch Bronconeumol.* 2021;57:739–40.
- Vaira LA, Salzano G, Le Bon SD, Maglio A, Petrocelli M, Steffens Y, et al. Prevalence of persistent olfactory disorders in patients with COVID-19: a psychophysical case-control study with 1-year follow-up. *Otolaryngol Head Neck Surg.* 2021;194:5998211061511.
- Han X, Fan Y, Alwalid O, Li N, Jia X, Yuan M, et al. Six-month follow-up chest CT findings after severe COVID-19 pneumonia. *Radiology.* 2021;299:e177–86.
- Clavario P, De Marzo V, Lotti R, Barbara C, Porcile A, Russo C, et al. Cardiopulmonary exercise testing in COVID-19 patients at 3 months follow-up. *Int J Cardiol.* 2021;340:113–8.
- Faverio P, Luppi F, Rebora P, Busnelli S, Stainer A, Catalano M, et al. Six-month pulmonary impairment after severe COVID-19: a prospective, multicentre follow-up study. *Respiration.* 2021;100:1078–87.
- Yin X, Xi X, Min X, Feng Z, Li B, Cai W, et al. Long-term chest CT follow-up in COVID-19 survivors: 102–361 days after onset. *Ann Transl Med.* 2021;9:1231.
- Zhao Y, Wang D, Mei N, Yin B, Li X, Zheng Y, et al. Longitudinal radiological findings in patients with COVID-19 with different severities: from onset to long-term follow-up after discharge. *Front Med.* 2021;8:711435.
- Bonazza F, Borghi L, di San Marco EC, Piscopo K, Bai F, Monforte AD, et al. Psychological outcomes after hospitalization for COVID-19: data from a multidisciplinary follow-up screening program for recovered patients. *Res Psychother.* 2020;23:491.
- Lombardo MDM, Foppiani A, Peretti GM, Mangiavini L, Battezzati A, Bertoli S, et al. Long-term coronavirus disease 2019 complications in inpatients and outpatients: a one-year follow-up cohort study. *Open Forum Infect Dis.* 2021;8:ofab384.
- Huang S, Zhou Z, Yang D, Zhao W, Zeng M, Xie X, et al. Persistent white matter changes in recovered COVID-19 patients at the 1-year follow-up. *Brain.* 2021; awab435.
- Du HW, Fang SF, Wu SR, Chen XL, Chen JN, Zhang YX, et al. Six-month follow-up of functional status in discharged patients with coronavirus disease 2019. *BMC Infect Dis.* 2021;21:1271.
- Fogante M, Cavagna E, Rinaldi G. COVID-19 follow-up: Chest X-ray findings with clinical and radiological relationship three months after recovery. *Radiography.* 2021;S1078–8174:00172–173.
- Wu Q, Zhong L, Li H, Guo J, Li Y, Hou X, et al. A follow-up study of lung function and chest computed tomography at 6 months after discharge in patients with coronavirus disease 2019. *Can Respir J.* 2021;2021:6692409.
- Stadnytskyi V, Bax CE, Bax A, Anfinrud P. The airborne lifetime of small speech droplets and their potential importance in SARS-CoV-2 transmission. *Proc Natl Acad Sci U S A.* 2020;117:11875–7.
- Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020;382:1708–20.
- Redd WD, Zhou JC, Hathorn KE, McCarty TR, Bazarbashi AN, Thompson CC, et al. Prevalence and characteristics of gastrointestinal symptoms in patients with severe acute respiratory syndrome coronavirus 2 infection in the United States: a multicenter cohort study. *Gastroenterology.* 2020;159:765–7, e762.
- Romero-Sánchez CM, Díaz-Maroto I, Fernández-Díaz E, Sánchez-Larsen A, Layos-Romero A, García-García J, et al. Neurologic manifestations in hospitalized patients with COVID-19: The ALBACOVID registry. *Neurology.* 2020;95:e1060–70.
- Lazarian G, Quinquenel A, Bellal M, Siavellis J, Jacquy C, Re D, et al. Autoimmune haemolytic anaemia associated with COVID-19 infection. *Br J Haematol.* 2020;190:29–31.
- Zulfiqar AA, Lorenzo-Villalba N, Hassler P, Andr C. Immune thrombocytopenic purpura in a patient with COVID-19. *N Engl J Med.* 2020;382:e43.
- Toscano G, Palmerini F, Ravaglia S, Ruiz L, Invernizzi P, Cuzzoni MG, et al. Guillain-Barré syndrome associated with SARS-CoV-2. *N Engl J Med.* 2020;382:2574–6.
- Feldstein LR, Rose EB, Horwitz SM, Collins JP, Newhams MM, Son MBF, et al. Multisystem inflammatory syndrome in U.S. children and adolescents. *N Engl J Med.* 2020;383:334–46.
- Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis.* 2020;20:425–34.
- Mo X, Jian W, Su Z, Chen M, Peng H, Peng P, et al. Abnormal pulmonary function in COVID-19 patients at time of hospital discharge. *Eur Respir J.* 2020;55:2001217.
- Gao Y, Chen R, Geng Q, Mo X, Zhan C, Jian W, et al. Cardiopulmonary exercise testing might be helpful for interpretation of impaired pulmonary function in recovered COVID-19 patients. *Eur Respir J.* 2021;57:2004265.
- Xie J, Covassin N, Fan Z, Singh P, Gao W, Li G, et al. Association between hypoxemia and mortality in patients with COVID-19. *Mayo Clin Proc.* 2020;95:1138–47.
- Mao R, Qiu Y, He JS, Tan JY, Li XH, Liang J, et al. Manifestations and prognosis of gastrointestinal and liver involvement in patients with COVID-19: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol.* 2020;5:667–78.
- Feng Z, Yu Q, Yao S, Luo L, Zhou W, Mao X, et al. Early prediction of disease progression in COVID-19 pneumonia patients with chest CT and clinical characteristics. *Nat Commun.* 2020;11:4968.
- Huang W, Li M, Luo G, Wu X, Su B, Zhao L, et al. The inflammatory factors associated with disease severity to predict COVID-19 progression. *J Immunol.* 2021;206:1597–608.
- Truffaut L, Demey L, Bruyneel AV, Roman A, Alard S, De Vos N, et al. Post-discharge critical COVID-19 lung function related to severity of radiologic lung involvement at admission. *Respir Res.* 2021;22:29.
- Mahase E. COVID-19: What do we know about “long COVID”? *BMJ.* 2020;370:m2815.
- Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. A clinical case definition of post-COVID-19 condition by a Delphi consensus. *Lancet Infect Dis.* 2022;22:e102–7.
- Guidelines NIffHaCEC. *COVID-19 rapid guideline: managing the long-term effects of COVID-19.* London: National Institute for Health and Care Excellence (UK); 2020.
- Groff D, Sun A, Ssentongo AE, Ba DM, Parsons N, Poudev GR, et al. Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review. *JAMA Netw Open.* 2021;4:e2128568.
- Nalbandian A, Sehgal K, Gupta A, Madhavan MV, McGroder C, Stevens JS, et al. Post-acute COVID-19 syndrome. *Nat Med.* 2021;27:601–15.
- Nasserie T, Hittle M, Goodman SN. Assessment of the frequency and variety of persistent symptoms among patients with COVID-19: a systematic review. *JAMA Netw Open.* 2021;4:e2111417.
- Fernández-de-Las-Peñas C, Palacios-Ceña D, Gómez-Mayordomo V, Florencio LL, Cuadrado ML, Plaza-Manzano G, et al. Prevalence of post-COVID-19 symptoms in hospitalized and non-hospitalized COVID-19 survivors: A systematic review and meta-analysis. *Eur J Intern Med.* 2021;92:55–70.
- Petrocelli M, Cutrupi S, Salzano G, Maglitto F, Salzano FA, Lechien JR, et al. Six-month smell and taste recovery rates in coronavirus disease 2019 patients: a prospective psychophysical study. *J Laryngol Otol.* 2021;135:436–41.
- de Lorenzo R, Cinel E, Cilla M, Compagnone N, Ferrante M, Falbo E, et al. Physical and psychological sequelae at three months after acute illness in COVID-19 survivors. *Panminerva Med.* 2021 Jun 1, <http://dx.doi.org/10.23736/S0031-0808.21.04399-8>. Online ahead of print.
- Chen Y, Ding C, Yu L, Guo W, Feng X, Yu L, et al. One-year follow-up of chest CT findings in patients after SARS-CoV-2 infection. *BMC Med.* 2021;19:191.
- Pan F, Yang L, Liang B, Ye T, Li L, Li L, et al. Chest CT patterns from diagnosis to 1 year of follow-up in COVID-19. *Radiology.* 2021;211199.
- Daher A, Balfanz P, Cornelissen C, Müller A, Bergs I, Marx N, et al. Follow up of patients with severe coronavirus disease 2019 (COVID-19): Pulmonary and extrapulmonary disease sequelae. *Respir Med.* 2020;174:106197.
- Steinbeis F, Thibeault C, Doellinger F, Ring RM, Mittermaier M, Ruwwe-Glösenkamp C, et al. Severity of respiratory failure and computed chest tomography in acute COVID-19 correlates with pulmonary function and respiratory symptoms after infection with SARS-CoV-2: An observational longitudinal study over 12 months. *Respir Med.* 2022;191:106709.
- Chen M, Liu J, Peng J, Jian W, Gao Y, Fang L, et al. Dynamic changes of pulmonary diffusion capacity in survivors of non-critical COVID-19 during the first six months. *EClinicalMedicine.* 2022;43:101255.
- Sonnweber T, Sahanic S, Pizzini A, Luger A, Schwabl C, Sonnweber B, et al. Cardiopulmonary recovery after COVID-19: an observational prospective multicentre trial. *Eur Respir J.* 2021;57.
- Desai AD, Lavelle M, Boursiquot BC, Wan EY. Long-term complications of COVID-19. *Am J Physiol Cell Physiol.* 2022;322:c1–11.
- Ramadan MS, Bertolino L, Zampino R, Durante-Mangoni E. Cardiac sequelae after coronavirus disease 2019 recovery: a systematic review. *Clin Microbiol Infect.* 2021;27:1250–61.
- Chan L, Chaudhary K, Saha A, Chauhan K, Vaid A, Zhao S, et al. AKI in hospitalized patients with COVID-19. *J Am Soc Nephrol.* 2021;32:151–60.
- Heung M, Steffick DE, Zivin K, Gillespie BW, Banerjee T, Hsu CY, et al. Acute kidney injury recovery pattern and subsequent risk of CKD: an analysis of veterans health administration data. *Am J Kid Dis.* 2016;67:742–52.
- Morin L, Savale L, Pham T, Colle R, Figueiredo S, Harrois A, et al. Four-month clinical status of a cohort of patients after hospitalization for COVID-19. *JAMA.* 2021;325:1525–34.
- Bangash MN, Patel J, Parekh D. COVID-19 and the liver: little cause for concern. *Lancet Gastroenterol Hepatol.* 2020;5:529–30.
- Guo H, Zhang Z, Zhang Y, Liu Y, Wang J, Qian Z, et al. Analysis of liver injury factors in 332 patients with COVID-19 in Shanghai, China. *Aging.* 2020;12:18844–52.
- Khoo B, Tan T, Clarke SA, Mills EG, Patel B, Modi M, et al. Thyroid function before, during, and after COVID-19. *J Clin Endocrinol Metabol.* 2021;106:e803–11.
- Higgins V, Sohaei D, Diamandis EP, Prassas I. COVID-19: from an acute to chronic disease? Potential long-term health consequences. *Crit Rev Clin Lab Sci.* 2021;58:297–310.
- Muñoz-Rodríguez JR, Gómez-Romero FJ, Pérez-Ortiz JM, López-Juárez P, Santiago JL, Serrano-Oviedo L, et al. Characteristics and risk factors associated with mortality in a multicenter Spanish cohort of patients with COVID-19 pneumonia. *Arch Bronconeumol.* 2021;57:34–41.
- Signes-Costa J, Núñez-Gil JJ, Soriano JB, Arroyo-Espiguelo R, Eid CM, Romero R, et al. Prevalence and 30-day mortality in hospitalized patients with COVID-19 and prior lung diseases. *Arch Bronconeumol.* 2021;57:13–20.
- Posso M, Comas M, Román M, Domingo L, Louro J, González C, et al. Comorbidities and mortality in patients with COVID-19 aged 60 years and older in a university hospital in Spain. *Arch Bronconeumol (Engl Ed).* 2020;56:756–8.

61. Tessitore E, Handgraaf S, Poncet A, Achard M, Höfer S, Carballo S, et al. Symptoms and quality of life at 1-year follow up of patients discharged after an acute COVID-19 episode. *Swiss Med Week*. 2021;151:w30093.
62. Seeße J, Waterboer T, Hippchen T, Simon J, Kirchner M, Lim A, et al. Persistent symptoms in adult patients one year after COVID-19: a prospective cohort study. *Clin Infect Dis*. 2021 Jul 5;ciab611, <http://dx.doi.org/10.1093/cid/ciab611>. Online ahead of print.
63. Caruso D, Guido G, Zerunian M, Polidori T, Lucertini E, Pucciarelli F, et al. Post-acute sequelae of COVID-19 pneumonia: six-month chest CT follow-up. *Radiology*. 2021;301:E396–405.
64. Fang X, Ming C, Cen Y, Lin H, Zhan K, Yang S, et al. Post-sequelae one year after hospital discharge among older COVID-19 patients: A multi-center prospective cohort study. *J Infect*. 2022;84:179–86.
65. Fumagalli C, Zocchi C, Tassetti L, Silverii MV, Amato C, Livi L, et al. Factors associated with persistence of symptoms 1 year after COVID-19: A longitudinal, prospective phone-based interview follow-up cohort study. *Eur J Intern Med*. 2022;97:36–41.
66. Heightman M, Prashar J, Hillman TE, Marks M, Livingston R, Ridsdale HA, et al. Post-COVID-19 assessment in a specialist clinical service: a 12-month, single-centre, prospective study in 1325 individuals. *BMJ Open Respir Res*. 2021;8.
67. Zhang X, Wang F, Shen Y, Zhang X, Cen Y, Wang B, et al. Symptoms and health outcomes among survivors of COVID-19 infection 1 year after discharge from hospitals in Wuhan, China. *JAMA Netw Open*. 2021;4:e2127403.
68. Crook H, Raza S, Nowell J, Young M, Edison P. Long covid-mechanisms, risk factors, and management. *BMJ*. 2021;374:n1648.
69. Karaarslan F, Güneri FD, Kardeş S. Long COVID: rheumatologic/musculoskeletal symptoms in hospitalized COVID-19 survivors at 3 and 6 months. *Clin Rheumatol*. 2022;41:289–96.
70. Taquet M, Dercon Q, Luciano S, Geddes JR, Husain M, Harrison PJ. Incidence, co-occurrence, and evolution of long-COVID features: A 6-month retrospective cohort study of 273,618 survivors of COVID-19. *PLoS Med*. 2021;18:e1003773.
71. Fernández-de-Las-Peñas C, Rodríguez-Jiménez J, Fuensalida-Novo S, Palacios-Ceña M, Gómez-Mayordomo V, Florencio LL, et al. Myalgia as a symptom at hospital admission by severe acute respiratory syndrome coronavirus 2 infection is associated with persistent musculoskeletal pain as long-term post-COVID sequelae: a case-control study. *Pain*. 2021;162:2832–40.
72. Maestre Muñiz MM, Arias Á, Mata-Vázquez E, Martín-Toledano M, López-Laramona G, Ruiz-Chicote AM, et al. Long-term outcomes of patients with coronavirus disease 2019 at one year after hospital discharge. *J Clin Med*. 2021;10.
73. Mattioli F, Piva S, Stampatori C, Righetti F, Mega I, Peli E, et al. Neurologic and cognitive sequelae after SARS-CoV2 infection: Different impairment for ICU patients. *J Neurol Sci*. 2022;432:120061.
74. Pilotto A, Cristillo V, Cotti Piccinelli S, Zoppi N, Bonzi G, Sattin D, et al. Long-term neurological manifestations of COVID-19: prevalence and predictive factors. *Neuro Sci*. 2021;42:4903–7.
75. Tleyjeh IM, Saddik B, AlSwaidan N, AlAnazi A, Ramakrishnan RK, Alhazmi D, et al. Prevalence and predictors of post-acute COVID-19 syndrome (PACS) after hospital discharge: A cohort study with 4 months median follow-up. *PLoS One*. 2021;16:e0260568.
76. Mazza MG, Palladini M, De Lorenzo R, Bravi B, Poletti S, Furlan R, et al. One-year mental health outcomes in a cohort of COVID-19 survivors. *J Psych Res*. 2021;145:118–24.
77. Zhou F, Tao M, Shang L, Liu Y, Pan G, Jin Y, et al. Assessment of sequelae of COVID-19 nearly 1 year after diagnosis. *Front Med*. 2021;8:717194.
78. González J, Benítez ID, Carmona P, Santisteban S, Monge A, Moncusí-Moix A, et al. Pulmonary function and radiologic features in survivors of critical COVID-19: a 3-month prospective cohort. *Chest*. 2021;160:187–98.
79. Wu X, Liu X, Zhou Y, Yu H, Li R, Zhan Q, et al. 3-month, 6-month, 9-month, and 12-month respiratory outcomes in patients following COVID-19-related hospitalisation: a prospective study. *Lancet Respir Med*. 2021;9:747–54.
80. Sanchez-Ramirez DC, Normand K, Zhaojun Y, Torres-Castro R. Long-term impact of COVID-19: a systematic review of the literature and meta-analysis. *BioMedicine*. 2021;9.
81. Xu B, Ma FQ, He C, Wu ZQ, Fan CY, Mao HR, et al. Incidence and affecting factors of pulmonary diffusing capacity impairment with COVID-19 survivors 18 months after discharge in Wuhan, China. *J Infect*. 2022;84:e16–8.
82. Montefusco L, Ben Nasr M, D'Addio F, Loretelli C, Rossi A, Pastore I, et al. Acute and long-term disruption of glycometabolic control after SARS-CoV-2 infection. *Nat Metab*. 2021;3:774–85.
83. Horwitz LI, Garry K, Prete AM, Sharma S, Mendoza F, Kahan T, et al. Six-month outcomes in patients hospitalized with severe COVID-19. *J Gen Intern Med*. 2021;36:3772–7.
84. Fernández-de-Las-Peñas C, Palacios-Ceña D, Gómez-Mayordomo V, Palacios-Ceña M, Rodríguez-Jiménez J, de-la-Llave-Rincón AI, et al. Fatigue and dyspnoea as main persistent post-COVID-19 symptoms in previously hospitalized patients: related functional limitations and disability. *Respiration*. 2022;101:132–41.
85. Xiong L, Li Q, Cao X, Xiong H, Huang M, Yang F, et al. Dynamic changes of functional fitness, antibodies to SARS-CoV-2 and immunological indicators within 1 year after discharge in Chinese health care workers with severe COVID-19: a cohort study. *BMC Med*. 2021;19:163.
86. Guillou A, Laurent E, Godillon L, Kimmoun A, Grammatico-Guillon L. Long-term mortality of elderly patients after intensive care unit admission for COVID-19. *Intensive Care Med*. 2021;47:710–2.
87. Munch MW, Granholm A, Nørregaard KM, Aksnes TS, Sølling CG, Christensen S, Perner A. Long-term mortality and health-related quality of life in the COVID STEROID trial. *Acta Anaesthesiol Scand*. 2022 Jan 24, <http://dx.doi.org/10.1111/aas.14029>.
88. Di Bari M, Tonarelli F, Balzi D, Giordano A, Ungar A, Baldasseroni S, et al. COVID-19, vulnerability, and long-term mortality in hospitalized and nonhospitalized older persons. *J Am Med Dir Assoc*. 2022;23:414–20.
89. Qin W, Chen S, Zhang Y, Dong F, Zhang Z, Hu B, et al. Diffusion capacity abnormalities for carbon monoxide in patients with COVID-19 at 3-month follow-up. *Eur Respir J*. 2021;58.
90. Cicco S, Vacca A, Cittadini A, Marra AM. Long-term follow-up may be useful in coronavirus disease 2019 survivors to prevent chronic complications. *Infect Chemother*. 2020;52:407–9.
91. Sun J, Liu J, Li H, Shang C, Li T, Ji W, et al. Pulmonary rehabilitation focusing on the regulation of respiratory movement can improve prognosis of severe patients with COVID-19. *Ann Pall Med*. 2021;10:4262–72.
92. Gloeckl R, Leitl D, Jarosch I, Schneeberger T, Nell C, Stenzel N, et al. Pulmonary rehabilitation in long COVID: more than just natural recovery? *ERJ Open Res*. 2021;7.
93. He Z, Ren L, Yang J, Guo L, Feng L, Ma C, et al. Seroprevalence and humoral immune durability of anti-SARS-CoV-2 antibodies in Wuhan, China: a longitudinal, population-level, cross-sectional study. *Lancet*. 2021;397:1075–84.
94. To KK, Hung IF, Ip JD, Chu AW, Chan WM, Tam AR, et al. Coronavirus Disease 2019 (COVID-19) re-infection by a phylogenetically distinct severe acute respiratory syndrome coronavirus 2 strain confirmed by whole genome sequencing. *Clin Infect Dis*. 2021;73:e2946–51.
95. Maier HE, Kuan G, Saborio S, Bustos Carrillo FA, Plazaola M, et al. Clinical spectrum of SARS-CoV-2 infection and protection from symptomatic re-infection. *Clin Infect Dis*. 2021 Aug 19;ciab717, <http://dx.doi.org/10.1093/cid/ciab717>. Online ahead of print.
96. Tan CW, Chia WN, Young BE, Zhu F, Lim BL, Sia WR, et al. Pan-sarbecovirus neutralizing antibodies in BNT162B2-immunized SARS-CoV-1 survivors. *N Engl J Med*. 2021;385:1401–6.
97. Killerby ME, Ata Ur, Rasheed M, Tamin A, Harcourt JL, Abedi GR, et al. Shedding of culturable virus, seroconversion, and 6-month follow-up antibody responses in the first 14 confirmed cases of coronavirus disease 2019 in the United States. *J Infect Dis*. 2021;224:771–6.
98. Wu X, Wang Z, He Z, Li Y, Wu Y, Wang H, et al. A follow-up study shows that recovered patients with re-positive PCR test in Wuhan may not be infectious. *BMC Med*. 2021;19:77.
99. Butterly S, Philip KEJ, Williams P, Fallas A, West B, Cumella A, et al. Patient symptoms and experience following COVID-19: results from a UK-wide survey. *BMJ Open Respir Res*. 2021;8.
100. Eloy P, Tardivon C, Martin-Blondel G, Isnard M, Turnier PL, Marechal ML, et al. Severity of self-reported symptoms and psychological burden 6-months after hospital admission for COVID-19: a prospective cohort study. *Int J Infect Dis*. 2021;112:247–53.
101. Hossain MA, Hossain KMA, Saunders K, Uddin Z, Walton LM, Raigangar V, et al. Prevalence of long COVID symptoms in Bangladesh: a prospective Inception Cohort Study of COVID-19 survivors. *BMJ Glob Health*. 2021;6.
102. Nesan G, Keerthana D, Yamini R, Jain T, Kumar D, Eashwar A, et al. 3-month symptom-based ambidirectional follow-up study among recovered COVID-19 patients from a tertiary care hospital using telehealth in Chennai, India. *Inquiry*. 2021;58, 469580211060165.
103. Xiong Q, Xu M, Li J, Liu Y, Zhang J, Xu Y, et al. Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study. *Clin Microbiol Infect*. 2021;27:89–95.
104. Bottemanne H, Gouraud C, Hulot JS, Blanchard A, Ranque B, Lahliou-Laforêt K, et al. Do anxiety and depression predict persistent physical symptoms after a severe COVID-19 episode? A prospective study. *Front Psych*. 2021;12:757685.
105. Augustin M, Schommers P, Stecher M, Dewald F, Gieselmann L, Gruell H, et al. Post-COVID syndrome in non-hospitalised patients with COVID-19: a longitudinal prospective cohort study. *Lancet Reg Health Eur*. 2021;6:100122.
106. Fernández-de-Las-Peñas C, Palacios-Ceña M, Gómez-Mayordomo V, Rodríguez-Jiménez J, Palacios-Ceña M, Velasco-Arribas M, et al. Long-term post-COVID symptoms and associated risk factors in previously hospitalized patients: A multicenter study. *J Infect*. 2021;83:237–79.
107. Chen X, Li Y, Shao TR, Yang LL, Li SJ, Wang XJ, et al. Some characteristics of clinical sequelae of COVID-19 survivors from Wuhan, China: A multi-center longitudinal study. *Influ Other Respir Virus*. 2021 Nov;19, 10.1111/irv.12943. doi: 10.1111/irv.12943. Online ahead of print.
108. Willi S, Lüthold R, Hunt A, Hänggi NV, Sejdju D, Scaff C, et al. COVID-19 sequelae in adults aged less than 50 years: A systematic review. *Travel Med Infect Dis*. 2021;40:101995.