

## Clinical Letter


### Wheelchair-assisted Ventilation in Neuromuscular Disease: The Abdominal Press-and-release Technique

Once respiratory muscle impairment has become pronounced, patients with neuromuscular disorders (NMD) can use non-invasive ventilation (NIV) to reduce symptoms related to CO<sub>2</sub> retention, improve functional status and quality of life (QoL) [1]. As respiratory conditions worsen, patients may require more extensive ventilatory support and transition from nighttime only to full time ventilator use, usually when experiencing daytime respiratory discomfort [2]. For some patients, nasal interface can be a comfortable option for daytime NIV, especially if they are able to tolerate it without skin breakdown. Other patients may use a nasal or oronasal interface for nighttime and switch to a mouthpiece for daytime use. In both cases, they experience limitations in their ability to perform daily activities, including communicating, eating or grooming. For this reason, finding a viable alternative to NIV to assist ventilation during daytime is a crucial point.

A 47-year-old female patient with severe Fukuyama congenital muscular dystrophy, a professor at our University, was evaluated

at the outpatient clinic of the Respiratory Pathophysiology Division of the University of Padua. She had been using home NIV via a nasal mask for 15 years prior to our clinical evaluation and regularly underwent lung recruitment manoeuvres through a mechanical in-exsufflator. She reported a history of progressive respiratory discomfort, fatigue and increasing need for ventilator use which had become necessary not only at night, but also during the day to alleviate symptoms. However, daytime use of NIV prevented her from giving lectures to her students. At that time, the patient realized that pressing her upper abdomen against the edge of the desk and releasing it rapidly by the wheelchair back and forth motion enhanced speech volume and duration, making it possible to perform uninterrupted lessons without ventilatory support; likewise, this technique enabled her to eat and drink without any assistance from the ventilator. As a result, she decided to use “wheelchair-assisted ventilation” during daytime whenever it was necessary to interrupt NIV. Remarkably, the patient discovered the method spontaneously and it was not a medical recommendation.

At our pulmonary function laboratory, the patient underwent pulmonary function test including recording of her pattern of breathing by a water-sealed bell spirometer (Baires 80, Biomedin,



**Fig. 1.** Breathing pattern recording and images obtained by electrical impedance tomography during unassisted breathing and wheelchair-assisted ventilation. Regional lung ventilation and the difference between the inspiratory and expiratory phase are shown in blue and grey color, respectively.

<https://doi.org/10.1016/j.arbres.2025.09.020>

0300-2896/© 2025 The Author(s). Published by Elsevier España, S.L.U. on behalf of SEPAR. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Padua, Italy) and electrical impedance tomography (EIT), a non-invasive monitoring tool providing continuous transverse images of the lungs, allowing a breath-to-breath evaluation of lung ventilation [3] (PulmoVista® 500, Dräger Medical GmbH, Lübeck, Germany). Her forced vital capacity (FVC) was found to be equal to 0.3 L. Arterial blood gas analysis showed normal values (PaO<sub>2</sub>: 78 mmHg; PaCO<sub>2</sub>: 44.5 mmHg; pH: 7.40; SaO<sub>2</sub>: 96%). After discontinuing NIV through oronasal mask, the patient could sustain unassisted breathing for about 90 s before the onset of respiratory discomfort. We found that tidal volume increased from 0.1 L during unassisted ventilation to 0.35 L during wheelchair-assisted ventilation at a respiratory rate of approximately 20 cycles/min (Fig. 1). Moreover, EIT showed a partial re-expansion of both lower lobes during the procedure (Fig. 1). SpO<sub>2</sub> value increased from 95% during unassisted breathing to 98% during assisted ventilation.

Similar to what happens during the application of the intermittent abdominal pressure ventilator [4], we hypothesized that compression of the upper abdomen against the edge of the desk could move the diaphragm upward to exhale air from the lungs by an increase in intra-abdominal pressure, whilst during release the diaphragm could return to its resting position due to gravity, thus fostering passive inspiration in combination with elastic return of the rib cage. As a result, we believe that wheelchair-assisted ventilation can effectively support breathing during daytime in NMD patients except for those cases with severe obesity, kyphoscoliosis, previous abdominal surgery, and/or gastroesophageal reflux. Required skills include ability to use commands for the electric wheelchair. Another limitation is that the user needs back extensors, and possibly some hip flexors to do the maneuver, both of which can be severely impaired due to underlying disease. Moreover, the technique cannot be effective in case of acute exacerbation, when patients need to use NIV 24/7 and mechanical in-exsufflation to treat oxygen desaturation. Since this is a single case and no systematic studies on long-term efficacy or safety are currently available, reproducibility and safety require further evaluation before the technique can be recommended. Despite limitations, wheelchair-assisted ventilation may represent a useful option to improve participation in daily activities and quality of life in selected NMD patients.

#### CRedit authorship contribution statement

Andrea Vianello: Conceptualization; writing – review & editing  
Gabriella Guarnieri: Investigation; original draft preparation, review & editing

Federico Lionello: Conceptualization, methodology, review & editing

#### Declaration of generative AI and AI-assisted technologies in the writing process

No content of the manuscript has been partially or totally produced with the help of any artificial intelligence software or tool.

#### Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

#### Conflict of interest

The authors declare not to have any conflicts of interest that may be considered to influence directly or indirectly the content of the manuscript.

#### References

- [1] AlBalawi MM, Castro-Codesal M, Featherstone R, Sebastianski M, Vandermeer B, Alkhaledi B, et al. Outcomes of long-term noninvasive ventilation use in children with neuromuscular disease: systematic review and meta-analysis. *Ann Am Thorac Soc* 2022;19:109–19, <http://dx.doi.org/10.1513/AnnalsATS.202009-1089OC>.
- [2] Racca F, Finder JD, Mercuri E, Vianello A, Chatwin M, Amaddeo A, et al. Current standards and future directions of Duchenne muscular dystrophy respiratory care: the PPMD Italy Meeting Report. *Pediatr Pulmonol* 2025;60, <http://dx.doi.org/10.1002/ppul.71113>, e71113.
- [3] Piraino T. An introduction to the clinical application and interpretation of electrical impedance tomography. *Respir Care* 2022;67:721–9, <http://dx.doi.org/10.4187/respcare.09949>.
- [4] Pierucci P, Di Lecce V, Carpagnano GE, Banfi P, Bach JR. The intermittent abdominal pressure ventilator as an alternative modality of noninvasive ventilatory support: a narrative review. *Am J Phys Med Rehabil* 2022;101:179–83, <http://dx.doi.org/10.1097/PHM.0000000000001804>.

Andrea Vianello\*, Gabriella Guarnieri, Federico Lionello

Department of Cardiac Thoracic Vascular Sciences and Public Health,  
University of Padova, Padova 35122, Italy

Corresponding author.

E-mail address: [andrea.vianello@aopd.veneto.it](mailto:andrea.vianello@aopd.veneto.it) (A. Vianello).