

5. Pierson DJ, Horton CA, Bates PW. Persistent bronchopleural air leak during mechanical ventilation. *Chest*. 1986;90:321–3. <http://dx.doi.org/10.1378/chest.90.3.321>.
6. Dugan KC, Laxmanan B, Murgu S, Hogarth K. Management of persistent air leaks. *Chest*. 2017;152:417–23. <http://dx.doi.org/10.1016/j.chest.2017.02.020>.
7. Carlon GC, Ray C, Klain M, McCormack P. Highfrequency positive-pressure ventilation in management of a patient with bronchopleural fistula. *Anesthesiology*. 1980;50:160–2.
8. Watanabe S, Watanabe T, Urayama H. Endobronchial occlusion method of bronchopleural fistula with metallic coils and glue. *Thorac Cardiovasc Surg*. 2003;51:106–8. <http://dx.doi.org/10.1055/s-2003-38981>.
9. Rodriguez Vial M, Lan C, Cornwell L, Omer S, Casal RF. Endobronchial oxygen insufflation: a novel technique for localization of occult bronchopleural fistulas. *Ann Am Thorac Soc*. 2013;10:157–9. <http://dx.doi.org/10.1513/AnnalsATS.201212-1260T>.
10. Loubriel B, Díaz J, Casal J, Rodriguez-Cintrón W. Sticking to life: closure of a bronchopleural fistula with fibrin glue in a patient with recurrent pneumothoraces. *J Bronchol*. 14:215–6. doi:10.1097/LBR.0b013e3180679e46.
11. Correia S, Dionisio J, Duro da Costa J. Modified technique of endobronchial balloon tamponade for persistent hemothorax. *J Bronchol Intervent Pulmonol*. 2014;21. <http://dx.doi.org/10.1097/LBR.000000000000102>.

Rodolfo Navarrete<sup>a,\*</sup>, Luis Pablo Cardonnet<sup>b</sup>,  
Lisandro Roberto Bettini<sup>b</sup>, Gonzalo Morosano<sup>c</sup>,  
Gustavo Cumbo Nacheli<sup>d</sup>

<sup>a</sup> Department of Pulmonary Medicine, Hospital Provincial del Centenario, Rosario, Argentina

<sup>b</sup> Department of Critical Care Medicine, Hospital Provincial del Centenario, Rosario, Argentina

<sup>c</sup> Pneumology, Argentina

<sup>d</sup> Bronchoscopy and Interventional Pulmonology, Pulmonary, Critical Care Medicine Division, Spectrum Health Medical Group Associate Director, Pulmonary and Critical Care Fellowship Program – Michigan State University, United States

\* Corresponding author.

E-mail address: [navarretehpc@gmail.com](mailto:navarretehpc@gmail.com)

(R. Navarrete).

<https://doi.org/10.1016/j.arbres.2020.10.006>

0300-2896/ © 2020 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

## Can we Train the Immune System of Patients With Cystic Fibrosis?



### ¿Podemos entrenar el sistema inmunológico de los pacientes con Fibrosis Quística?

Dear Editor,

The importance of exercise in cystic fibrosis (CF) is well known since regular training is able to reduce lung function decline and to improve exercise tolerance and quality of life.<sup>1</sup> Moreover, exercise can enhance muco-ciliary clearance by inhibiting epithelial sodium channel (ENaC) conductance, decreasing Na re-absorption and favouring airway fluid secretion.<sup>2</sup> Therefore, current international guidelines recommend it as a physiotherapy adjunct to improve the efficacy of airway clearance techniques (ACTs).<sup>3</sup>

Beyond that, little is known about the effect of physical activity on inflammation and immune defence mechanisms in CF patients; only few studies are available, focusing mainly on pulmonary physiotherapy techniques.<sup>4</sup> Differently, some evidences are available in other diseases: in chronic obstructive pulmonary disease (COPD) training programs have demonstrated to exert a positive effect on lymphocyte proliferative response.<sup>5</sup> Studies on mice models showed that moderate exercise could induce a shift in immune response from a Th1 to a Th2 response.<sup>6</sup> This change seemed to be associated with reduced viral or bacterial load, inflammation, infection-related morbidity and mortality.<sup>6,7</sup>

We conducted a prospective interventional single-centre pilot study in the Cystic Fibrosis Unit of our Hospital from February to July 2019, to investigate whether an exercise program can exert a modulator effect on immune system of adults with CF and chronic bronchial infection (CBI).

Inclusion criteria were: stable adult CF patients with CBI, body mass index (BMI) > 19, forced expiratory volume in one second (FEV<sub>1</sub>) between 30% and 90% of predicted value, exclusion of immunological or autoimmune disorders. Written informed consent to participate in the study was obtained from participants. Ethics approval was granted by the Ethics Committee of our Hospital.

Exercise training program: A 12-week individualized training program in a dedicated gym under the supervision of a professional personal trainer was planned, with 4 sessions per week of


45–60 min each (adapted to individual physical capability), focused on cardiovascular resistance and strength. A minimum of 3 sessions/week was required to patients. Individual starting capacity for the exercise training program was based on the evaluation of physical fitness during the first week of training and the exercise test at first visit. Patients were instructed how to perform the exercises at the start of the study and their progression was evaluated by the personal trainer once a week. The personal routine was mostly organized in order to train between 60% and 75% of maximal initial effort, based on baseline exercise test, and to have 50% of training dedicated to cardiovascular resistance and 50% to strength, but personalization of the training program was set by the trainer according to individual characteristics.

Within 7 days before the training, protocol included clinical and functional assessment, complete lung function tests (forced spirometry, diffusion capacity of the lungs for carbon monoxide –DLCO– and plethysmography), cardiopulmonary exercise test (CPET), blood and sputum collection. The same procedures were scheduled one week after the end of the training in order to avoid the effects of acute immune response.

For statistical analysis, data normality was checked using Kolmogorov–Smirnov and Pearson omnibus normality tests. For data comparison, Paired *t* test was used to compare parametric data, and the Wilcoxon–Mann–Whitney test was used to compare non-parametric data. Significance was set at *P* < 0.05. GraphPad Prism 8.0 package (GraphPad Software, Inc, USA) was used.

Six patients were recruited, two female and four males, aged 20–33 years (mean 26.17 ± 5.23), all with class I or II CFTR mutations and 3 with pancreatic insufficiency (F508del was present in 5 out of 6 patients, being homozygosis in only 1 patient). Four patients were on chronic treatment with fluticasone (100 to 500mcg/daily) in combination with salmeterol, while 2 patients only used salbutamol on demand. Mean FEV<sub>1</sub> was 57.2 ± 23.6% of predicted, DLCO/VA was 97.7 ± 17.7%, Total Lung Capacity (TLC) 104.2 ± 10.5%.

CBI was due to methicillin-sensitive *Staphylococcus aureus* (MSSA) and *Pseudomonas aeruginosa* (PA) in 5/6 and 4/6 patients respectively. Three patients had both MSSA and PA chronic infection. The target for completion of the training was of 4 sessions per week during the 12 weeks of the study (100%). A specific questionnaire demonstrated a mean completion of the training of



**Fig. 1.** The figure represents only the variables showing significant changes between T<sub>0</sub> (baseline) and T<sub>1</sub> (end of the study). (1.1) Parameters of cardiopulmonary test at T<sub>0</sub> and T<sub>1</sub> measure points. Five patients are represented in the figure since one patient (FQ-AF-003) did not achieve maximal effort exercise at T<sub>1</sub>. (A) The ventilatory equivalent for carbon dioxide (EqCO<sub>2</sub>) significantly decreased while (B) the Ventilatory reserve (BR) significantly increased. (1.2) Percentages of T and B cells in peripheral blood samples before and after the training period for each subject. Blood lymphocytes subpopulations are represented: (C) A significant decrease in the percentage of CD4+ effector memory (TEM) and central memory (TCM) T cells at T<sub>1</sub> was observed. Th17 subpopulation (D), recent thymic emigrant cells population (RET) (E) and B CD4+ naive subpopulation (F) showed a mild increase at the end of the training period (T<sub>1</sub>). (1.3) Results of sputum cell count at T<sub>0</sub> and T<sub>1</sub>. Neutrophils and macrophages significantly predominated over eosinophils and lymphocytes at both T<sub>0</sub> and T<sub>1</sub>. P values for T<sub>1</sub> -vs- T<sub>0</sub> comparisons are represented as follows: \* *p* < 0.03; \*\* *p* = 0.002; \*\*\*\* *p* < 0.0001.

74 ± 10.25% (standard deviation). Only one patient poorly followed the program due to fact he doubled his workload in a car service achieving similar training effects; therefore he was maintained in the study for observational purpose. Analysis of advanced peripheral blood immunophenotype was performed by flow cytometry.<sup>8</sup> At the same time, spontaneous (5 patients) or induced (1 patient) sputum was collected and processed for differential leukocyte count by microscopy.

One patient with chronic MSSA and PA infection had a mild exacerbation at week 10 of training, treated with amoxicillin/clavulanate for 8 days. During that period her training programme was suspended. This complication did not impact on her FEV1 or on her capacity of exercise; unfortunately, it was not possible to analyse blood or sputum during exacerbation.

At the end of the training period, lung function tests did not show significant changes.

In comparison with baseline, post-training CPET showed a significant decrease in ventilatory equivalent for carbon dioxide (EqCO<sub>2</sub>) from 30.1% to 28.7% ( $p < 0.0001$ ), while breathing reserve (BR) increased significantly in all subjects (10.2% to 24.5%,  $p = 0.002$ ) (Fig. 1.1).

In peripheral blood samples, no significant changes were found in the main lymphocyte subpopulations. However, a significant decrease in the percentage of CD4+ effector memory (TEM) and central memory (TCM) T cells was observed after training (44.0–38.16%,  $p < 0.03$ ). The analysis of T-helper cell functional subsets showed a significant percentage increase in Th17 subset cells (17.2–23.9%;  $p < 0.03$ ), while Th1 and Th2 tended to decrease (37–30.6%;  $p = 0.18$  and 33.6–20.3,  $p = 0.19$  respectively) and Th1-17 showed a non-significant increase (14.5–25.2%;  $p = 0.10$ ). Recent thymic emigrant cells (RTEs) significantly increased in all subjects (36.9–40.9%;  $p < 0.03$ ) such as the B naive subpopulation (IgD + CD27–) from 66.4% to 68.6% ( $p < 0.03$ ) (Fig. 1.2).

Sputum cell differential count showed significant predominance of neutrophils and macrophages over other cell types in both phases of the study ( $p < 0.0001$ ). Compared to baseline, no significant differences were detected after the training ( $p = 0.99$ ), but mild reduction in neutrophils was observed (–7%, Fig. 1.3).

In conclusion, as expected, our pilot study confirms that training was able to improve efficacy of ventilation by increasing BR and decreasing EqCO<sub>2</sub>.

Beyond that, we also found interesting changes in blood lymphocytes subpopulation after training. A significant increase of the Th17 cells was observed in peripheral blood; these cells play a role in adaptive immunity against pathogens, especially PA,<sup>9</sup> maintaining mucosal barriers and contributing to pathogen clearance.<sup>10</sup> In particular, there is clear evidence about the role of Th17 cells in modulating both systemic and airways inflammation in CF.<sup>11,12</sup> On the other hand, the observed increase of RTEs may indicate a moderate expansion of the T-naive cell repertoire (CD45RA+ CCR7+) (that in fact increased in 4 patients), which represents a positive effect on the immune response,<sup>13</sup> while the reduction of TEM and TCM may favour a better control of the local infection.

Lastly, the mild reduction in neutrophils observed in sputum suggests that exercise might be capable of reducing neutrophilic inflammation by improving airways clearance and ventilatory efficacy.

This pilot study has several limitations, including: limited sample size, lack of a control group, small number of measures, absence of absolute cellular count.

Despite that, our preliminary trial may suggest that also the immune system is positively influenced by exercise training. This new research hypothesis deserves in our opinion further investigation particularly by increasing sample size, in order to finally improve CF outcomes in the future.

## Conflict of interests

Eva Polverino has received consultancy and speaker's fees from: Bayer, Grifols, Insmad, Chiesi, Menarini, Zambon, Pfizer, Teva, Shire and Polyphor. The remaining authors have no conflicts of interest.

## Acknowledgements

The authors sincerely thank Prof. R Pujol Borrell for his advice and review of the manuscript.

## References

1. Hebestreit H, Kriemler S, Radtke T. Exercise for all cystic fibrosis patients: is the evidence strengthening? *Curr Opin Pulm Med*. 2015.
2. Schmitt L, Wiebel M, Frese F, Dehnert C, Zugck C, Bärtsch P, et al. Exercise reduces airway sodium ion reabsorption in cystic fibrosis but not in exercise asthma. *Eur Respir J*. 2011.
3. Flume PA, Robinson KA, O'Sullivan BP, Finder JD, Vender RL, Willey-Courand D-B, et al. Cystic fibrosis pulmonary guidelines: airway clearance therapies. *Respir Care*. 2009.
4. Moeller A, Stämpfli SF, Rueckert B, Rechsteiner T, Hamacher J, Wildhaber JH. Effects of a short-term rehabilitation program on airway inflammation in children with cystic fibrosis. *Pediatr Pulmonol*. 2010;45:541–51.
5. Fernandes JR, Marques da Silva CCB, da Silva AG, de Carvalho Pinto RM, da Silva Duarte AJ, Carvalho CR, et al. Effect of an Exercise Program on Lymphocyte Proliferative Responses of COPD Patients. *Lung*. 2018.
6. Lowder T, Padgett DA, Woods JA. Moderate exercise early after influenza virus infection reduces the Th1 inflammatory response in lungs of mice. *Exerc Immunol Rev*. 2006;12:97–111.
7. Stravinskias Durigon T, Mackenzie BA, Carneiro Oliveira-Junior M, Santos-Dias A, De Angelis K, Malfitano C, et al. Aerobic exercise protects from *Pseudomonas aeruginosa*-induced pneumonia in elderly mice. *J Innate Immun*. 2018.
8. Garcia-Prat M, Álvarez-Sierra D, Aguiló-Cucurull A, Salgado-Perandrés S, Briongos-Sebastian S, Franco-Jarava C, et al. Extended immunophenotyping reference values in a healthy pediatric population. *Cytom B – Clin Cytom*. 2019.
9. Muntaka S, Almuhan Y, Jackson D, Singh S, Afryie-Asante A, Cámara M, et al. Gamma interferon and interleukin-17A differentially influence the response of human macrophages and neutrophils to *pseudomonas aeruginosa* infection. *Infect Immun*. 2019.
10. Singh B, Schwartz JA, Sandrock C, Bellemore SM, Nikoipour E. Modulation of autoimmune diseases by interleukin (IL)-17 producing regulatory T helper (Th17) cells. *Indian J Med Res*. 2013.
11. Tan HL, Regamey N, Brown S, Bush A, Lloyd CM, Davies JC. The Th17 pathway in cystic fibrosis lung disease. *Am J Respir Crit Care Med*. 2011;184:252–8.
12. Kushwah R, Gagnon S, Sweezey NB. Intrinsic predisposition of naïve cystic fibrosis T cells to differentiate towards a Th17 phenotype. *Respir Res*. 2013;14:138.
13. McFarland RD, Douek DC, Koup RA, Picker LJ. Identification of a human recent thymic emigrant phenotype. *Proc Natl Acad Sci USA*. 2000.

Jordi Tur-Torres<sup>a,1</sup>, Letizia Traversi<sup>b,1</sup>, Mónica Martínez-Gallo<sup>c,d</sup>, Giulio Assante<sup>e</sup>, Christian Eduardo Romero Mesones<sup>a</sup>, David Clofent Alarcon<sup>a,b,c,d,e,f</sup>, Patricia Chang-Macchiu<sup>a</sup>, Antoni Alvarez<sup>f</sup>, Eva Polverino<sup>a,b,c,d,e,f,\*</sup>

<sup>a</sup> Institut de Recerca Vall d'Hebron (VHIR), Hospital Universitari Vall d'Hebron (HUVH), Universitat Autònoma de Barcelona, Spain

<sup>b</sup> Department of Medicine and Surgery, Respiratory Diseases, Università dell'Insubria, Varese-Como, Italy

<sup>c</sup> Immunology Division, Department of Clinical and Molecular Genetics, Hospital Universitari Vall d'Hebron (HUVH), Vall d'Hebron Research Institute (VHIR), Barcelona, Spain

<sup>d</sup> Department of Cell Biology, Physiology and Immunology, Universitat Autònoma de Barcelona (UAB), Barcelona, Spain

<sup>e</sup> Personal trainer DIR, Barcelona, Spain

<sup>f</sup> Pneumology Department, Vall d'Hebron University Hospital (HUVH), Vall d'Hebron Research Institute (VHIR), CIBER, Barcelona, Spain

\* Corresponding author.

E-mail address: [eva.polverino@vhir.org](mailto:eva.polverino@vhir.org) (E. Polverino).

<sup>1</sup> These authors equally contributed to this manuscript as first authors.

<https://doi.org/10.1016/j.arbres.2020.10.014>

0300-2896/© 2020 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.