



Letter to the Director

Swimming Exercise for Patients With Long-Term Respiratory Post COVID-19 Complications: Further Thinking on the Pulmonary Rehabilitation



To the Director,

Swimming is an aquatic sport in which the propulsion is coordinated with respiratory phases, consisting in underwater apnoeas interspersed with rapid and deep inhalations out of the water. This breathing pattern provokes those aquatic athletes have higher lung volumes and lung diffusing capacities than other athletes and general population.¹

The SARS-coronavirus-2 disease 19 (COVID-19) has put the respiratory health in the spotlight. Some factors, as advanced age, and obesity has been correlated with a higher risk of hospitalization and mortality. On the other hand, exercise capacity has been reported as a positive factor to reduce the clinical impact of COVID-19 infection.

The benefits of physical exercise over human being health have been extensively described. However, the structural and functional properties of the lungs do not change significantly due to training, except for aquatic athletes. Swimming involves both locomotor muscles as respiratory muscles, which may be combined to offer a complete pulmonary fitness program. Why do not translate these benefits to the pulmonary rehabilitation (PR) programs based in physical exercise?

Three months after hospital discharge, patients with COVID-19 still present lung function abnormalities with a 25–57% of the patients showing carbon monoxide lung diffusion capacity (DL_{CO}) < 80% of reference, confirming that survivors of COVID-19 may suffer lung function sequelae.²

For patients with COVID-19, PR is aimed at relieving symptoms of dyspnoea, psychological distress and improving the functional status. Exercise training is considered the foundation of PR and it is included in most of the rehabilitation programs. Some of the current guidelines relative to management of COVID-19 convalescent people only propose walking or cycling activities, being the potential benefits of swimming still misunderstood, unknown, and lacking research.

Conventional PR programmes rely on general principles of exercise physiology: (duration, intensity, and frequency), but

the impact of exercise modality is not considered. Some specific attributes of swimming may stimulate thoracic expansion and increased gas diffusing capacity, such as horizontal aerobic exercise, water hydrostatic pressure, and forced submaximal inspirations.³

Our hypothesis is that different exercise modalities may trigger different rehabilitation results, based on the biomechanical and physiological attributes of each locomotion mode. At this moment, swimming exercise has not been utilized as a PR modality, in comparison with the classical rehabilitation methods, such as respiratory muscle training and walking exercise. Some studies on COPD patients have used water-based exercise, but only focused on a combination of strength exercises and mobility in the water (*aqua-gym*).⁴

As far as we know, no studies have been conducted using swimming as a rehabilitation exercise method. The main reason could be the difficulty of getting access to a swimming pool. Also, swimming biomechanics is more difficult to master than walking or cycling, and it demands a multidisciplinary work between medical doctors and exercise trainers. Besides, some patients may feel a discomfort feeling with their face underwater since patients recovering from COVID-19 present respiratory complications with augmented exercise ventilation.⁵

However, especially for those with previous experience, swimming exercise may be a valuable rehabilitation method. Future studies must explore this paradigm, comparing different exercise modalities to evaluate the efficacy of the existing PR programs.

References

- García I, Drobnic F, Pons V, Viscor G. Acute changes in lung diffusing capacity after training in elite swimmers. *Arch Bronconeumol*. 2021;57:306–7.
- Sibila O, Albacar N, Perea L, Faner R, Torralba Y, Hernandez-Gonzalez F, et al. Lung Function sequelae in COVID-19 patients 3 months after hospital discharge. *Arch Bronconeumol*. 2021;57:59–61.
- García I, Drobnic F, Arrillaga B, Pons V, Viscor G. Lung capacity and alveolar gas diffusion in aquatic athletes: implications for performance and health. *Apunts Sports Med*. 2021:56.
- McNamara RJ, McKeough ZJ, McKenzie DK, Alison JA. Water-based exercise in COPD with physical comorbidities: a randomised controlled trial. *Eur Respir J*. 2013;41:1284–91.
- Baratto C, Caravita S, Faini A, Perego GB, Senni M, Badano LP, et al. Impact of COVID-19 on exercise pathophysiology: a combined cardiopulmonary and echocardiographic exercise study. *J Appl Physiol*. 2021;130:1470–8.

Iker García^{a,*}, María Molina-Molina^b, Beatriz Arrillaga^c,
Casimiro Javierre^d, Ginés Viscor^a

^a Secció de Fisiologia, Departament de Biologia Cel·lular, Fisiologia i Immunologia, Facultat de Biologia, Universitat de Barcelona, Av. Diagonal, 643, 08028 Barcelona, Spain

^b Unidad Funcional de Intersticio Pulmonar (UFIP), Servicio de Neumología, Hospital Universitario de Bellvitge, IDIBELL, CIBERES, Hospitalet de Llobregat 08907, Spain

^c Unit of Human Anatomy and Embryology, Department of Pathology and Experimental Therapeutics, Faculty of Medicine and Health Sciences, University of Barcelona, 08907 Hospitalet de Llobregat, Spain

^d Faculty of Medicine and Health Sciences, University of Barcelona, 08907 Hospitalet de Llobregat, Spain

Corresponding author.

E-mail address: ikergarciaalday@gmail.com (I. García).