



Letters to the Editor

Promotion of anti-smoking strategies as the most effective and efficient way to reduce lung cancer (and other diseases)*

Fomento de las estrategias antitabaco como la manera más efectiva y eficiente de reducir el cáncer de pulmón (y otras enfermedades)

To the Editor,

We read with interest the article entitled, “Estimation of the risk of lung cancer in women participating in a population-based breast cancer screening program”, published by González-Marrón et al., and we would like to highlight some issues with this paper.¹ First, the authors assume that the results of the NELSON study are valid, although these had not been published in full at the time of submission of this letter. The only results available are from the presentation made by the lead author of the study at the World Conference on Lung Cancer in 2018, where he pointed out that the NELSON study does not have sufficient power in women, since they accounted for only 16% of the sample (the author states: “includes a small subgroup of women and, the trial was initially powered for high risk males”). Accordingly, we believe it is arguable that the results of the NELSON study are applicable to women in this study.²

Second, the title of the article itself is confusing. Strictly speaking, the study presented does not make a risk estimate, but quantifies the percentage of women who would meet the screening criteria proposed by the NLST and NELSON studies, respectively. Equating “risk estimation” with the fulfillment of screening criteria could confuse readers, especially when numerous recently published studies have suggested that risk estimation in lung cancer (LC) screening, if performed, should be individualized for each patient, as the criteria used by both studies, particularly the NLST, imply a high percentage of false positives.

Third, LC screening continues to be the subject of debate in Europe and the United States.³ It is remarkable that the possibility of screening in Spain is being suggested when experts cannot even agree on the periodicity of this screening. The NELSON and NLST studies have different screening schedules, and the NELSON study also has rounds of varying duration. Few people know, moreover, that although the Medicare and Medicaid Committees indicated that the benefit/risk ratio of LC screening was unfavorable,⁴ LC screening was introduced in the United States following pressure from senators and congressmen.

Finally, the authors should have highlighted the enormous opportunity cost of funding smoking cessation treatments compared to the implementation of screening, as has been shown in Spain.⁵ This funding would prevent respiratory, cardiovascular, and numerous types of cancer, significantly increase life expectancy in these new ex-smokers and, in addition, involves far less iatrogenesis than participation in an LC screening program. In our opinion, more emphasis should have been placed on the need to promote strategies such as the World Health Organization’s MPOWER measures (<https://www.who.int/tobacco/mpower/en/>) to reduce tobacco use in Spain, particularly among women who can attend any population screening program, not just for breast cancer.

References

- González-Marrón A, Martín-Sánchez JC, García-Alemany F, Martínez-Martín E, Matilla-Santander N, Cartanyà-Hueso A, et al. Estimación del riesgo de cáncer de pulmón en mujeres que participan en un programa de cribado poblacional de cáncer de mama. Arch Bronconeumol. 2019. <http://dx.doi.org/10.1016/j.arbres.2019.04.014>.
- De Koning HJ, Van Der Aalst CM, ten Haaf K, Oudkerk M. Available from: https://library.iaslc.org/conference-program?product_id=10&author=&category=&date=2018-09-25&session.type=&session=&presentation=&keyword=&available=&cme=&page=2, 2018.
- Verghese C, Redko C, Fink B. Screening for lung cancer has limited effectiveness globally and distracts from much needed efforts to reduce the critical worldwide prevalence of smoking and related morbidity and mortality. J Glob Oncol. 2018;4:1–7.
- Centers for Medicare and Medicaid Services. Available from: <https://www.cms.gov/medicare-coverage-database/details/medcac-meeting-details.aspx?MEDCACId=68>, 2014.
- Ruano-Ravina A, Provencio-Pulla M, Casan Clarà P. Lung cancer screening with low-dose computerised tomography. Thoughts on its use in Spain. Med Clin (Barc). 2016;147:366–70.

 Alberto Ruano-Ravina,^{a,b,*} Mariano Provencio-Pulla,^{c,d}
 José Alberto Fernández-Villar^e
^a Área de Medicina Preventiva y Salud Pública, Universidad de Santiago de Compostela, Santiago de Compostela, Spain

^b CIBER de Epidemiología y Salud Pública, CIBERESP, Madrid, Spain

^c Servicio de Oncología, Hospital Puerta de Hierro, Majadahonda, Madrid, Spain

^d Instituto de Investigación Puerta de Hierro, Majadahonda, Madrid, Spain

^e Grupo NeumoVigo I+i, Instituto de Investigación Sanitaria Galicia Sur (IISGS), Servicio de Neumología, Hospital Álvaro Cunqueiro, Vigo, Pontevedra, Spain

* Corresponding author.

 E-mail address: alberto.ruano@usc.es (A. Ruano-Ravina).

<https://doi.org/10.1016/j.arbr.2019.12.003>

1579-2129/ © 2020 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

* Please cite this article as: Ruano-Ravina A, Provencio-Pulla M, Fernández-Villar JA. Fomento de las estrategias antitabaco como la manera más efectiva y eficiente de reducir el cáncer de pulmón (y otras enfermedades). Arch Bronconeumol. 2020;56:339.